

CLAIMS

We claim:

1. A composition (for inhibition of inducible COX-2) activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof.

2. The composition of Claim 1 wherein first and second components are derived from plants or plant extracts.

3. The composition of Claim 1 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or disaccharides, amino acids, sulfates, succinate, acetate and glutathione.

4. The composition of Claim 1, formulated in a pharmaceutically acceptable carrier.

5. The composition of Claim 1, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

5 603
6. A composition for inhibition of inducible COX-2 activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and ^{terpene} helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, aneoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof.

10
7. The composition of Claim 6 wherein first and second components are derived from plants or plant extracts.

15 603
8. The composition of Claim 6 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or di- saccharides, amino acids, sulfates, succinate, acetate and glutathione.

9. The composition of Claim 6, formulated in a pharmaceutically acceptable carrier.

10. The composition of Claim 6, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

11. A composition for inhibition of inducible COX-2 activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof.

12. The composition of Claim 11 wherein first and second components are derived from plants or plant extracts.

13. The composition of Claim 11 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or di- saccharides, amino acids, sulfates, succinate, acetate and glutathione.

14. The composition of Claim 11, formulated in a pharmaceutically acceptable carrier.

15. The composition of Claim 11, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

16. A composition for inhibition of inducible COX-2 activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof.

17. The composition of Claim 16 wherein first and second components are derived from plants or plant extracts.

18. The composition of Claim 16 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or di- saccharides, amino acids, sulfates, succinate, acetate and glutathione.

19. The composition of Claim 16, formulated in a pharmaceutically acceptable carrier.

20. The composition of Claim 16, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

21. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

22. The method of Claim 21 wherein the composition is formulated in a dosage form such that said administration provides from 0.05 to 5.0 mg body weight per day of each sequesterpene lactone species, and from 0.5 to 20.0 mg/kg bodyweight per day of each diterpene lactone species or triterpene species.

23. The method of Claim 21, wherein the composition is administered in an amount sufficient to maintain a serum concentration of 0.001 to 10 μ M of each sesquiterpene lactone species and from 0.001 to 10 μ M of each diterpene lactone or triterpene species.

24. The method of Claim 21 wherein said animal is selected from the group consisting of humans, non-human primates, dogs, cats, birds, horses and ruminants.

25. The method of Claim 21 wherein administration is by a means selected from the group consisting of oral, parenteral, topical, transdermal and transmucosal delivery.

26. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, aneoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

27. The method of Claim 26 wherein the composition is formulated in a dosage form such that said administration provides from 0.05 to 5.0 mg body weight per day of each sequesterpene lactone species, and from 0.5 to 20.0 mg/kg bodyweight per day of each diterpene lactone species or triterpene species.

5

28. The method of Claim 26, wherein the composition is administered in an amount sufficient to maintain a serum concentration of 0.001 to 10 μ M of each sesquiterpene lactone species and from 0.001 to 10 μ M of each diterpene lactone or triterpene species.

29. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

30. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first

component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

31. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

32. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid,

triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

33. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

34. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

35. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

36. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

37. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the

group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

5

38. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

10

39. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

15

20

5
10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500
510
520
530
540
550
560
570
580
590
600
610
620
630
640
650
660
670
680
690
700
710
720
730
740
750
760
770
780
790
800
810
820
830
840
850
860
870
880
890
900
910
920
930
940
950
960
970
980
990
1000

40. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

41. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

42. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

5

09519506-073104